- **1.** Given $f(x) = \sin x + 2\cos x$, $-\frac{\pi}{4} \le x \le 0$, find $(f^{-1})'(a)$ for a = 2. Ans: $(f^{-1})'(2) = 1$ Note: $f^{-1}(2) = 0$
- **2.** Solve for x:
- (a). $e^{3x-7} = 6$ $x = \frac{7+\ln 6}{3}$
- **(b).** $\ln(x+1) \ln x = 1$ $x = \frac{1}{2 \cdot 1}$

- **3.** Given $f(x) = \ln(\cos x + 2)$
- (a). What is the domain of f? Domain: all real
- (b). Find the relative extreme values on the interval [-1,1].

[Indicate max or min.]

Max value of $\ln 3$ at x = 0

4. Evaluate the following integrals.

(a).
$$\int_0^1 \frac{1}{4 - 2x} dx = -\frac{1}{2} \ln|4 - 2x| \Big|_0^1 = \frac{1}{2} \ln 2$$

(d).
$$\int \sec(5x) dx = \frac{1}{5} \ln|\sec(5x) + \tan(5x)| + C$$

(b).
$$\int e^{-7x} - \frac{7 \ln x}{x} dx = -\frac{1}{7} e^{-7x} - \frac{7}{2} (\ln x)^2 + C$$

(e).
$$\int_0^{\ln 2} \frac{e^{3x} + 1}{e^x} dx = \frac{1}{2} e^{2x} - e^{-x} \Big|_0^2 = 2$$
Simplify using log./exp. properties.

(c).
$$\int \frac{3x^2 + 2x}{x^3 + x^2 + 1} dx = \ln|x^3 + x^2 + 1| + C$$

(f).
$$\int \frac{\tan x}{\ln(\cos x)} dx = -\ln|\ln(\cos x)| + C$$

- **5.** Given the function $f(x) = e^{-x^2}$
- (a). Evaluate $\lim_{x \to -\infty} e^{-x^2} = 0$ (b). Evaluate $\lim_{x \to +\infty} e^{-x^2} = 0$ (c). Find $f'(x) = -2xe^{-x^2}$

6. What is the formula for $\log_a x$ in terms of the natural logarithmic function?

 $\log_a x = \frac{\ln x}{\ln a}$

7. Differentiate the following functions:

(a).
$$y = x^4 - 4^x + e^{4x} + \ln 4x$$

 $y' = 4x^3 + 4^x \cdot \ln 4 + 4e^{4x} + \frac{1}{x}$

(c).
$$y = \pi^x - \ln e^x$$
 $y' = \pi^x \cdot \ln \pi - 1$

$$y' = \pi^x \cdot \ln \pi - 1$$

(b).
$$y = x^{\frac{1}{x}}$$

$$y' = \frac{1}{x^2} (1 - \ln x) \cdot x^{1/x}$$

(b).
$$y = x^{\frac{1}{x}}$$
 $y' = \frac{1}{x^2}(1 - \ln x) \cdot x^{1/x}$ (d). $h(\theta) = 3\ln\left(\frac{2 + \cos\theta}{\theta^2}\right)$ $h'(\theta) = \frac{-3\sin\theta}{2 + \cos\theta} - \frac{6}{\theta}$


$$h'(\theta) = \frac{-3\sin\theta}{2 + \cos\theta} - \frac{6}{\theta}$$

8. Find the equation of the tangent line to $y = \log_3 x$ at x = 1

$$y = \frac{1}{\ln 3}(x-1) .$$

9. (a). Find the maximum area of a rectangle in the first quadrant with 2 sides on the x- and y-axes and one vertex on the curve $y = e^{-x}$. See the figure below. [Hint: Express the area of such a rectangle in terms of x only.] Maximize A = xy subject to $y = e^{-x}$

 $\Longrightarrow x = 1$. So max area $= \frac{1}{a}$



- (b). Sketch the picture for a rectangle in the first quadrant with 2 sides on the x- and y-axes and one vertex on the curve $y = e^x$. Without using Calculus, determine whether there exists such a rectangle with a maximum area. Briefly explain (a couple of sentences) why or why not. No, if you draw the picture you will see that you can always create a bigger rectangle by taking x further out.
- 10. Given that a population follows the law of exponential growth, $y(t) = Ce^{kt}$ where y is the population and t is time in years.
- (a). Find the proportional constant k, if the population triples every 50 years.

$$k = \frac{\ln 3}{50}$$

(b). If the population is 100 after 5 years, find the population at time t.

$$y = \frac{100}{\frac{10}{3}} e^{\frac{\ln 3}{50}t} = 89.60e^{.02197t}$$

- 11. Find the following limits. Clearly show all steps and indicate where you use L'Hopital's Rule.

- (a). $\lim_{x\to 0} \frac{\sin 4x}{2x} = 2$ (b). $\lim_{x\to \infty} (1+x)^{\frac{1}{x}} = 1$ (c). $\lim_{x\to 2} \frac{x-2}{x^2-3x+2} = 1$ (d). $\lim_{x\to 2} \frac{x-2}{x^2-3x-2} = 0$