

[Give critical values $t_{\alpha/2}$ to 3 decimal places.]

1. Given a confidence level of 95%,

(a). Sketch a picture and determine the value of $\alpha/2$.

(b). Use `invT` to find the critical value $t_{\alpha/2}$ for a 95% confidence level and the following sample sizes.

(i) Sample size $n = 25$

$$t_{.025} = 2.064$$

(ii) Sample size $n = 50$

$$t_{.025} = 2.010$$

(iii) Sample size $n = 100$

$$t_{.025} = 1.984$$

(iv) Sample size $n = 200$

$$t_{.025} = 1.972$$

2. Given a confidence level of 90%,

(a). Sketch a picture and determine the value of $\alpha/2$.

(b). Use `invT` to find the critical value $t_{\alpha/2}$ for a 90% confidence level and the following sample sizes.

(i) Sample size $n = 48$

$$t_{.05} = 1.678$$

(ii) Sample size $n = 182$

$$t_{.05} = 1.653$$

3. Given a confidence level of 99%,

(a). Sketch a picture and determine the value of $\alpha/2$.

(b). Use `invT` to find the critical value $t_{\alpha/2}$ for a 99% confidence level and the following sample sizes.

(i) Sample size $n = 64$

$$t_{.005} = 2.656$$

(ii) Sample size $n = 256$

$$t_{.005} = 2.595$$