[Note: The Final Exam is comprehensive. Use the old review sheets, exams, and quizzes to study previous material.]

1. Evaluate the following integrals. [**Note:** You may or may not need to use substitution.] Check your answer by differentiating the result.

(a)
$$\int_{0}^{2} t^{2} \sqrt{1 + t^{3}} dt = \frac{52}{9}$$
 assubstitution
(b)
$$\int \sin x \cos(\cos x) dx = -\sin(\cos x) + C$$
 assubstitution
(c)
$$\int 3x^{5} - 4x^{3} + 6x + 2 dx = \frac{1}{2}x^{6} - x^{4} + 3x^{2} + 2x + C$$
 direct integration
(d)
$$\int (3x - 1)(3x^{2} - 2x)^{2} dx = \frac{1}{6}(3x^{2} - 2x)^{3} + C$$
 assubstitution
(e)
$$\int x(3x^{2} - 2x)^{2} dx = \frac{3}{2}x^{6} - \frac{12}{5}x^{5} + x^{4} + C$$
 expand/simplify
(f)
$$\int \left(1 + \frac{1}{t}\right) \left(\frac{1}{t^{2}}\right) dt = -\frac{1}{t} - \frac{1}{2t^{2}} + C$$
 multiply/simplify OR assubstitution
(g)
$$\int_{0}^{\pi/6} \sec x \tan x dx = \frac{2}{\sqrt{3}} - 1$$
 direct integration rule
(h)
$$\int \sin x \cos x dx = \frac{1}{2} \sin^{2} x + C \text{ OR } - \frac{1}{2} \sin^{2} x + C$$
 assubstitution
(j)
$$\int_{1}^{5} \frac{5x}{\sqrt{1 - x^{2}}} dx = \frac{15}{4}(1 - x^{2})^{2/3} + C$$
 assubstitution
(j)
$$\int_{1}^{3} \frac{x^{2} + 1}{x^{2}} dx = \frac{8}{3}$$
 simplify
(k)
$$\int y^{3} \sqrt{y} dy = \frac{2}{7}y^{7/2} + C$$
 simplify
(l)
$$\int_{0}^{1} (2 - x)^{6} dx = \frac{127}{7}$$
 assubstitution
(m)
$$\int \theta \sin(3\theta^{2}) d\theta = -\frac{1}{6}\cos(3\theta^{2}) + C$$

(a).
$$f(x) = 3 - 2x - x^2$$
, $g(x) = -x + 1$ $\frac{9}{2}$ (b). $x = y^2$, $x = -y$ $\frac{1}{6}$

3. Find the volume of the solid generated by rotating the region bounded by the given curves about the given line.

(a). $y = x^2, y = 4x - x^2$ about the line y = 6(b). xy = 6, y = 2, y = 6, x = 6 about the line x = 6. [Set up, but do not evaluate!!!] $V = \int_{2}^{6} \pi \left(6 - \frac{6}{y} \right)^{2} dy$

4. The force exerted by gravity on an object sent into space is given by $F(x) = \frac{4.8 \times 10^{11}}{x^2}$ pounds where x is measured in miles from the *center* of the earth. How much work is done to propel a satellite module to 800 miles above the earth. [Use 4000 miles for the radius of the earth]

 $W = \int_{4000}^{4800} 4.8 \times 10^{11} x^{-2} \, dx = 2 \times 10^7 \text{ mile} \cdot \text{lbs} = 1.056 \times 10^{11} \text{ foot} \cdot \text{lbs}$

5. If 18 J of work is required to stretch a spring 40 cm from it's natural length, find the work required to stretch it an additional 30 cm.

 $k = \frac{9}{400}$ (if using cm) k = 225 (if converted to m) $\Rightarrow W = 37.125$ J

6. Given $f(x) = \frac{4x^2 + 4}{x^2}$

(a). Find the average value of f(x) on the interval [1,3].

(b). Use the Mean Value Theorem for integrals to find all values x = c where $f(c) = f_{ave}$. $x = \sqrt{3}$

7. Find the value of k so that the average value of $f(x) = kx^2 - x$ on [0,2] is equal to 4. $k = \frac{15}{4}$

 $f_{ave} = \frac{16}{3}$