1. Given the following information, find the values of the remaining trigonometric functions.

$$\tan\theta = 3, \quad \pi < \theta < \frac{3\pi}{2}. \qquad \qquad \sin\theta = -\frac{3}{\sqrt{10}}, \ \cos\theta = -\frac{1}{\sqrt{10}}, \ \tan\theta = 3, \ \csc\theta = -\frac{\sqrt{10}}{3}, \ \sec\theta = -\sqrt{10}, \ \cot\theta = \frac{1}{3}$$

2. Solve the following equations for x.

(a).
$$2\sin^2 x - \sqrt{2}\sin x = 0$$
 (x in $[0, 2\pi]$) $x = 0, \pi, 2\pi, \frac{\pi}{4}, \frac{3\pi}{4}$ (b). $\cos\left(\frac{x}{2}\right) = 0$ (x in $[0, 2\pi]$) $x = \pi$

3. Given
$$\theta = \frac{3\pi}{4}$$
, find $\sin 2\theta$.

4. Differentiate the following using *Differentiation Rules*.

(a).
$$y = 10x^3 - 3x + 7$$

(b).
$$f(x) = \pi^2$$

(c).
$$y = (3x)^3$$
 $\frac{dy}{dx} = 81x^2$

(d).
$$y = \frac{x + 4x^3 - 3}{x^3}$$
 $y' = \frac{x^3 (1 + 12x^2) - (x + 4x^3 - 3)(3x^2)}{x^6}$ or $y' = -2x^{-3} + 9x^{-4}$

(e).
$$s(t) = t^2(3t - 4t^3)$$

(f).
$$f(x) = \frac{3}{x^2} - \sqrt{x}$$

(g).
$$s(t) = (3t^3 - t^2 + 7)^{23}$$

$$s'(t) = 23(3t^3 - t^2 + 7)^{22}(9t^2 - 2t)$$

(h).
$$f(\theta) = \theta \sin(\theta^2 + 1)$$
 $f'(\theta) = 2\theta^2 \cos(\theta^2 + 1) + \sin(\theta^2 + 1)$

(i).
$$y = \frac{x(2x^4 + 4)^8}{\tan 2x}$$
 [Do not simplify!]
$$\frac{dy}{dx} = \frac{\tan 2x \cdot \left[x \cdot 8(2x^4 + 4)^7 \cdot 8x^3 + (2x^4 + 4)^8 \cdot 1\right] - x(2x^4 + 4)^8(\sec^2(2x) \cdot 2)}{\tan^2 2x}$$

5. Find the equation of the tangent line to the curve
$$y = \sqrt[3]{2x^2 - 5}$$
 at $x = 4$. $y - 3 = \frac{16}{27}(x - 4)$

6. Given
$$f(x) = g(3x^2)$$
, find f' in terms of g' .
$$f'(x) = g'(3x^2) \cdot 6x$$

7. A tank holds 1000 gallons of water, which drains from the bottom of the tank in 50 minutes. Torricelli's Law gives the volume V of water remaining in the tank after t minutes as $V = 1000 \left(1 - \frac{1}{50}t\right)^2$ for $0 \le t \le 50$. Find the rate at which the water is draining from the tank after 10 minutes. Include units in your answer.

-32 gallons/min.

8. If a stone is thrown vertically upward on the moon with a velocity of 8 m/s, its height after t seconds is given by $y = 8t - 0.83t^2$, [Calculator*]

- (a). What is the velocity after 2 s?
- $4.68 \mathrm{m/s}$
- **(b).** What is the velocity at impact?
- -8 m/s

- **9.** The cost function for a certain commodity is $C(x) = 60 + 0.12x 0.0004x^2 + .000002x^3$.
 - $C'(x) = 0.12 0.0008 x + 0.000006 x^2$

(a). Find the marginal cost function.

 $\mathcal{F}(x) = 0.12 - 0.0008 \, x + 0.000006 \, x^2$

(b). Find and interpret C'(50).

C'(50) = \$0.095.

[Calculator*]

The rate the cost is changing when the 50th item is produced is approximately \$0.095 per item.

(c). Compare C'(50) with the cost of producing the 51st item.

C(51) - C(50) = \$0.094902

10. Any Section 2.7 applications.

11. Given the curve drawn below and defined by


$$x^2 + y^2 = 3 + xy$$

(a). Find $\frac{dy}{dx}$

$$\frac{dy}{dx} = \frac{y - 2x}{2y - x}$$

- (b). On the graph below, sketch any tangents lines to the curve where the slope is 0.
- (c). Use part (a) to find these points on the curve where the slope is 0. Must show work for credit. (1,2) & (-1,-2).
- (d). Find $\frac{d^2y}{dx^2}$ in terms of x and y.

$$\frac{d^2y}{dx^2} = \frac{(2y-x)\left(\frac{y-2x}{2y-x}-2\right) - (y-2x)\left(2\frac{y-2x}{2y-x}-1\right)}{(2y-x)^2}$$



[The problem below is from Section 2.9, which will be covered on Tuesday.]

- **12.** Given $f(x) = \sqrt{x} = x^{1/2}$
- (a). Find the linearization L(x) at a=25

$$L(x) = 5 + \frac{1}{10}(x - 25)$$

- (b). Use this linearization L(x) to approximate $\sqrt{24.7}$
- [Simplify your answer.]

4.97

(c). Find the differential dy for x going from 25 to 25.5.

0.05

^{*}Similar non-calculator problems could be given.