

Before we can prove this, we need more theorems.

PROOF Let  $E$  be an  $n \times n$  elementary matrix.

(a). For row replacement, there are only 2 possible elementary matrices:

(c). For row scaling by  $k$ , there are only 2 possible elementary matrices:

i.e., it is true for  $n = k + 1$ . Therefore, by mathematical induction it is true for all  $n \times n$  elementary matrices. ■

THEOREM If  $A$  is an  $n \times n$  matrix and  $E$  is an  $n \times n$  elementary matrix, then

$$\det(EA) = \det(E) \det(A) = \begin{cases} \det(A), & \text{if row replacement} \\ -\det(A), & \text{if row interchange} \\ k \det(A), & \text{if row scaled by } k \end{cases}$$

PROOF Do As Homework. [Hint: Similar to last proof.]

Note: This is Theorem 3 on p. 169, with  $B$  written as  $EA$ .

EX:  $A = \begin{bmatrix} 2 & 4 \\ 1 & -5 \end{bmatrix}$   $\det A =$

$-2R_2 + R_1 \rightarrow R_1 :$   $B_1 =$

$R_2 \leftrightarrow R_1 :$   $B_2 =$

$-3R_2 \rightarrow R_2 :$   $B_3 =$

THEOREM If  $A$  is an  $n \times n$  matrix that is reduced to echelon form  $U$  using **only** row interchanges and row replacements. Then

$$\begin{aligned} \det A &= (-1)^r \det U && \text{where } r \text{ is the number of row interchanges used.} \\ &= (-1)^r && \text{(i.e. } \underline{\hspace{2cm}} \text{ )} \\ &&& \text{since the echelon form is always upper triangular.} \end{aligned}$$

EX: Reduce the following matrix to echelon form and compute the determinant. Compare with computing the determinant by cofactor expansion.

$$A = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 3 & -2 \\ 0 & 1 & 2 \end{bmatrix}$$